

**Huskie for Kochia Control in Winter Wheat - 2007**  
 Robert Klein and Jeffrey Golus, University of Nebraska, North Platte  
 Kevin Watteyne, Bayer CropScience

Location: WCREC Dryland Farm, North Platte.

Procedure: Plots were laid out in randomized complete block design with three replications. Kochia was seeded into standing winter wheat in April, but stand was inadequate to place a trial. Treatments were then applied to kochia in fallow with no winter wheat present. Treatments were applied June 21, 2007 with a ten foot boom (six XR11002 nozzles on 20 inch spacing) at 13.5 gpa carrier volume and 22 psi nozzle pressure. Kochia was 1-3 inches tall. Plots were rated visually for percent control July 3 and July 14.

Treatment	Rate Prod/a	Timing	Kochia Percent Control	
			July 3	July 14
1. Untreated Check			0	0
2. Huskie + NIS + AMS	11 fl oz + 0.5% + 0.5 lb	Spring	69	69
3. Huskie + NIS + UAN	11 fl oz + 0.5% + 2 qt	Spring	53	63
4. Huskie + NIS + UAN	15 fl oz + 0.5% + 2 qt	Spring	74	84
5. Huskie + Ally + NIS + UAN	11.3 fl oz + 0.1 oz + 0.5% + 2 qt	Spring	100	100
LSD 5%			7.2	7.2